An Assessment of Cheyenne River Sioux Tribe Watersheds Impaired by Nonpoint Source Pollution Mitra N. Jha, Ph.D. Ecosystem Protection Program US EPA Region-8, Denver, CO 80202

Summary:

On the Chevenne River Sioux Tribe (CRST) Reservation most Nonpoint Source (NPS) Pollution is caused by agriculture, Hydromodification, land disposal, and resource extraction. The non-discrete nature of NPS pollution presents significant difficulties for CRST in reference to monitoring and managing surface water resources. Best Management Practice pilots have been started to reduce or control the NPS pollution. The Structural and Nonstructural controls and operational and maintenance procedures have been applied to reduce the NPS pollution activities

NPS pollution Characteristics



Occurs over extensive areas Enters receiving water in a diffuse manner Carried by water over land or through soil profile to receiving water Difficult to trace to point of origin Magnitude related to climatic events

Indian Reservations in Region 8

27 Tribes and 26 Reservations Region 8 Indian Reservation land: 28,468, 062 acres where as US Indian Reservation land: 69,420,257 acres 41% of land located in Region-8 with approximately 60.500 stream miles

Socioeconomic Condition of CRST

West Central South Dakota- 4.230 miles 70% Indian & 30% non-Indian Population Agriculture is the primary activity Unemployment-70% Farmers and Ranchers supplement income through off-farm employment

Chevenne River Sioux Tribes Land Base: (2,787,207 acres)

(Trust and allotted-1,391,58998, Feeland-1,300,193, and total water areas-95,424 acres)

Land Use:

Rangeland- 81% Cropland- 18% Others 1%

(including Forest land, Low density urban, Construction, Small water areas etc)

Chevenne River ST Reservation in South Dakota



Beneficial/Use Categories

Domestic Water Supply Fish and aquatic and Wildlife propagation Irrigation and agricultural supply Tribal Cultural and Spiritual Recreation



Surface Water Quality


Fifty eight Streams- drain to the Missouri River Sixteen Watersheds group established Two watersheds sampled each year since 2001 Quality of surface water ranges from moderate to very poor depending on water source and time of year Water quality is better in spring and poor in summer and fall

Source of Impairment at CRST

Known-9 water bodies, unknown-59



Agriculture number one industry 767 farms responsible for 90% NPS, Nutrients-fertilizers, wastes, natural etc-nuisance growth of algae, weeds, Misuse of pesticides, decrease dissolved oxygen level, Increase total dissolved solids, sediments and Mining activities- Metals, Other pollutants



Degree of NPS Impacts at CRST

7 Water bodies- Severe impacts 41 Water bodies- Moderate impacts 16 Water bodies- Slight impacts Assessment Level:

-9 Water bodies monitored -55 Water bodies evaluated

CRST Best Management Practices

General Management Program:

-Administration, assessment, monitoring, and general education

Specific Management Program:

-BMPs associated with agriculture, land disposal, hydro-modification, and resource extraction

CRST Pilot Project:

Location: Rousseau Creek Challenges: Stream bank/channel alteration, Construction etc.

Selected a drainage in southeastern part Rousseau Creek- 25 miles, impaired, riparian 16 producers with different agricultural habitat

Solution: Structural and nonstructural control, sediment

Vegetation filter strips-Replanting naturally occurring vegetation (cotton woods, willows, sedges, and legumes) in several hundred square feet or smaller. Other BMPs site specific for each agriculture and land disposal activities, hydro-modification



Conclusion:

NPS pollution is larger problem at CRST than available resources. Tribes need more scientific, technical, and financial support to solve the water quality problems

